

5
**METHODS AND COMPOSITIONS FOR ENHANCING CELLULAR FUNCTION
THROUGH PROTECTION OF TISSUE COMPONENTS**

10
Cross-Reference to Related Applications

This application claims the benefit of provisional application number 60/200,843 filed May 1, 2000; provisional application number 60/233,263 filed September 6, 2000; and provisional application number 60/233,025 filed September 15, 2000; each entitled "Methods and Compositions for Enhancing Cellular Function and Protecting Receptor."

15
Background of the Invention

Cellular function depends on the maintenance of intact cellular components including: receptors, proteins, lipids, nucleic acids, carbohydrates, hormones and cofactors. Cellular receptors, including cell surface receptors, mediate communication within and between cells, tissues and organs within a living system. Cellular receptors also provide a means to signal a living system, tissues, organs, cells, and subcellular compartments. Receptors are molecules or macromolecules that bind or interact with agents to alter or enhance their function. Many receptors are membrane bound proteins, which require not only that their protein structure be intact but also that the membrane lipids and carbohydrates be intact and functional. Through various signaling mechanisms, the messages sent by the receptor, either in the presence or absence of an interacting or bound agent, can be transmitted. Following receptor activation, signalling also requires intact cellular proteins, lipids, nucleic acids and carbohydrates in order for the message to be properly received.

20
25

Often as a result of damage, the ability of cellular receptors to interact with or bind various agents is decreased, resulting in an impairment of vital intrinsic and extrinsic communication. Damage to cellular receptors and other cellular components diminishes the ability of a receptor to bind agents and elicit a communication or signaling event. This can result in damage or death to cells, resulting in damage or diseases of tissues, organs and living systems. Accordingly, there is a need for a means to protect receptors and other cellular components from damage and to increase the efficacy of agents that exert their effects through cellular receptors.

Summary of the Invention

The invention provides methods for enhancing cellular function through protection of tissue components and/or increasing the efficacy of a therapeutic agent in a subject in need thereof. The method includes administering a composition, such as a pharmaceutical composition, of a pyrophosphate analog. In a second embodiment, the method includes administering a composition, such as a pharmaceutical composition, of a protective agent.


Preferably, the invention provides a method for protecting a muscarinic acetylcholine receptor (mAChR) and/or increasing the efficacy of an agent that directly or indirectly affects a mAChR in a subject in need thereof. Suitable agents that directly or indirectly affect a muscarinic receptor include anticholinesterase agents, muscarinic agonists, allosteric regulators of a muscarinic receptor, muscarinic antagonists, and neurotrophic and neuritogenic factors that are similar to naturally occurring nerve growth promoting substances. In one embodiment, the invention provides a method to protect a mAChR and/or increase the efficacy of agents that directly or indirectly affect a mAChR in the central

nervous system (CNS) of a subject in need thereof. Preferably, a muscarinic receptor is protected from an endogenous low molecular weight inhibitor from Alzheimer's brain tissue, a metal, or oxidative stress. In another embodiment, the invention provides a method to protect a mAChR and/or increase the efficacy of agents that directly or indirectly affect a
5 mAChR not in the CNS of a subject in a need thereof. In a first embodiment, the method includes administering a pyrophosphate analog. In a second embodiment, the method includes administering a protective agent.

The invention also provides a method for increasing the efficacy of a therapeutic agent, preferably a neurologic agent, in a subject in need thereof. In a first embodiment, this
10 method includes administering a pyrophosphate analog. In this embodiment, the increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or induced by the pyrophosphate analog. In a second embodiment, this method includes administering a protective agent. In this second embodiment, the increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or
15 induced by the protective agent. In each embodiment, the subject preferably is concurrently receiving, has recently received, or will soon receive a neurologic agent such as nerve growth factor (NGF), insulin growth factor (IGF-1), brain derived neurotrophic factor (BDNF), fibroblast growth factor (FGF), and the like; certain other known neurotrophins and neuroprotectants; and medications used for stroke, Alzheimer's disease, Parkinson's disease,
20 ALS, traumatic brain or spinal cord injury, cancer, diabetes, neuropathies, hypertension, irritable bowel syndrome; diseases or disorders of the heart and smooth muscles, blood, blood vessels, glands or bones and other disorders. Preferably, the therapeutic agent directly or


indirectly affects a mAChR. Such agents include an anticholinesterase agent, a muscarinic agonist, and a muscarinic antagonist.

Pyrophosphate analogs that can be employed in the appropriate embodiment of the method of the invention include compounds of Formula I:



where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²), and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900. Compounds of Formula I in which R¹ is a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²); or R² is a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl can be referred to as substituted pyrophosphate analogs. Compounds of Formula I can also include substituted pyrophosphate analogs such as a dinucleoside-5-5'-pyrophosphate, a cyclopyrophosphate of purine, and a pyrimidine acyclonucleoside. The compound of Formula I can be any pharmaceutically acceptable salt or basic addition salt. Preferably, X is O, CH₂, NH, or S; R¹ is H; R² is H; and n is 1-6. More preferably the pyrophosphate analog is pyrophosphate or imidodiphosphate.

Additional pyrophosphate analogs include compounds of Formula II:



where n=2-4; X is O, RCR¹; CR; C (n=4), CH (n=3), or CH₂ (n=2); NH; N; S; and R and/or R¹ is H, OH, a small alkyl group, such as CH₃ or (CH₂)_mNH₂ where m=1-6. Further included are bisphosphonic acids, which are also known as bisphosphonates, where X is preferably RCR¹, where R and R¹ groups are chosen independently from OH, H₂N(CH₂)₂, or CH₃. For example, RCR¹ can be H₂N(CH₂)₂C(OH) or CH₃COH. More specifically, the bisphosphonates include etidronic acid ((1-Hydroxyethylidene)bisphosphonic acid) and pamidronic acid ((3-Amino- 1-hydroxypropylidene)bisphosphonic acid) where preferably n=2.

Yet more additional pyrophosphate analogs include substituted pyrophosphate analogs such as an inositol diphosphate, an inositol triphosphate, an inositol tetraphosphate, an inositol pentaphosphate, and an inositol hexaphosphate.

Suitable protective agents that can be employed in an embodiment of the method of the invention include a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavinoid, a combination thereof, or a pharmaceutically acceptable salt thereof; a heme binding compound, such as hemopexin, lipopexin, a lipoprotein, or ApoE-2; and a heme oxygenase, such as heme oxygenase-1 or heme oxygenase-2, biliverdin reductase, a catalase, a peroxidase, a vector encoding a biliverdin reductase, a vector encoding a heme oxygenase (e.g. a vector encoding a heme oxygenase-1 or a vector encoding a heme oxygenase-2), a vector encoding a catalase, a vector encoding a peroxidase, or a combination thereof. Biliverdin reductase can be administered alone or in combination with a heme oxygenase.

Heme oxygenases include recombinant heme oxygenase. Preferably, a heme oxygenase is a human heme oxygenase.

The method of the invention can treat or prevent a CNS disorder. Preferably, the method of the invention can treat or prevent neurodegeneration, can improve memory and cognition, can treat or prevent brain deterioration or cognitive and memory loss associated with aging, or can treat or prevent Alzheimer's Disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, and/or schizophrenia; nerve damage from cerebrovascular disorders such as stroke or atherosclerosis in the brain or spinal cord, from CNS infections including meningitis and HIV, from tumors of the brain and spinal cord, prion diseases, and CNS disorders resulting from ordinary aging (e.g., anosmia), brain injury, or spinal cord injury.

In another embodiment, the method of the invention can treat or prevent a disease or disorder not of the CNS. Preferably, the method of the invention can treat or prevent cancer, or neuropathies or diseases or disorders of the heart, smooth muscles, blood, blood vessels, glands, or bones. Such diseases or disorders include hypertension, myocardial infarction, ischemic heart disease, congestive heart failure, cardiac arrhythmias, cancer, irritable bowel syndrome, diverticular disease, urinary incontinence, esophageal achalasia, chronic obstructive airways disease, xerostomia, diabetes mellitus, Sjogren's syndrome or dry eye syndrome which involves decreased secretion of tears by, for example, the lacrimal glands, Paget's disease, hereditary hematochromatosis or a non-CNS disorder resulting from normal aging.

In another embodiment, the method of the invention treats infections, including (without limitation) bacterial, fungal, algo, or algae infections. Such infections can occur in plants (for which a preferred embodiment of the invention employs imidodiphosphate as the pyrophosphate analog), animals, or mammals.

5

Brief Description of the Figures

Figure 1 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Pyrophosphate protected the receptor from loss of antagonist (^3H -QNB (quinulidinyl benzilate)) binding.

10

Figure 2 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate protected the mAChR from inactivation by heme and peroxide. Pyrophosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 3 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Pyrophosphate protected the receptor from loss of agonist (oxotremorine) binding.

15

Figure 4 illustrates protection of a mAChR by imidodiphosphate. Imidodiphosphate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Imidodiphosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

20

Figure 5 illustrates protection of a mAChR by guanylimidodiphosphate. Guanylimidodiphosphate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Guanylimidodiphosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 6 illustrates protection of a mAChR by adenylylimidodiphosphate.

Adenylylimidodiphosphate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Adenylylimidodiphosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

5 Figure 7 illustrates protection of a mAChR by tripolyphosphate. Tripolyphosphate protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Tripolyphosphate protected the receptor from loss of antagonist (^3H -QNB) binding.

10 Figure 8 illustrates protection of a mAChR by bilirubin. Bilirubin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Bilirubin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 9 illustrates protection of a mAChR by bilirubin. Bilirubin protected the mAChR from inactivation by heme and peroxide. Bilirubin protected the receptor from loss of antagonist (^3H -QNB) binding.

15 Figure 10 illustrates protection of a mAChR by bilirubin. Bilirubin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Bilirubin protected the receptor from loss of agonist (oxotremorine) binding.

Figure 11 illustrates protection of a mAChR by biliverdin. Biliverdin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Biliverdin protected the receptor from loss of antagonist (^3H -QNB) binding.

20 Figure 12 illustrates protection of a mAChR by carnosol, Carnosol protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Carnosol protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 13 illustrates protection of a mAChR by quercetin. Quercetin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Quercetin protected the receptor from loss of antagonist (^3H -QNB) binding.

5 Figure 14 illustrates protection of a mAChR by myricetin. Myricetin protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Myricetin protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 15 illustrates protection of a mAChR by catalase. Catalase protected the mAChR from inactivation by the endogenous low molecular weight inhibitor. Catalase protected the receptor from loss of antagonist (^3H -QNB) binding.

10 Figure 16 illustrates protection of a mAChR by catalase. Catalase protected the mAChR from inactivation by heme and peroxide. Catalase protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 17 illustrates protection of a mAChR by a peroxidase. The peroxidase protected the mAChR from inactivation by the endogenous low molecular weight inhibitor.
15 The peroxidase protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 18 illustrates protection of a mAChR by a peroxidase. The peroxidase protected the mAChR from inactivation by heme and peroxide. The peroxidase protected the receptor from loss of antagonist (^3H -QNB) binding.

Figure 19 illustrates protection of mAChR by pamidronate. Pamidronate protected the
20 mAChR from inactivation by the endogenous low molecular weight inhibitor.

Figure 20 illustrates protection of a mAChR by pyrophosphate. Pyrophosphate protected the mAChR from damage by the metal lead in the form of PbCl_2 . Pyrophosphate protected the receptor from loss of antagonist (^3H -QNB (quinulidinyI benzilate)) binding.

Detailed Description of the Invention

Definitions

5 As used herein, “cholinesterase” refers to an enzyme capable of hydrolyzing acetylcholine and includes acetylcholinesterase.

 As used herein, “agonist” refers to an agent that binds to or interacts with a receptor and elicits a response transduced through the receptor. Agonist includes full agonists, partial agonists, and inverse agonists. A full agonist is an agent that can elicit a maximal response
10 from a receptor. A partial agonist is an agent that can elicit, at best, a less than maximal response from a receptor. An inverse agonist is an agent that produces a response that is opposite that of a full or partial agonist. For example, if agonist binding or interaction with a receptor results in increased concentration of cAMP within a cell, then inverse agonist
15 binding or interaction with the same receptor will result in a decreased concentration of cAMP within the cell.

 As used herein, “antagonist” refers to an agent that is capable of partially or completely inhibiting, or reversing, the effect of an agonist on a receptor.

 As used herein, “allosteric modifier” refers to an agent that binds or interacts with a site other than the agonist binding site of a receptor and modifies the ability of an agonist or
20 an antagonist to elicit or inhibit, respectively, a response transduced through a receptor, without itself eliciting a response.

 As used herein, “tissue component” includes receptors, proteins, lipids, nucleic acids, carbohydrates, hormones, vitamins, and cofactors.

As used herein, "receptor" refers to any molecule or macromolecule within or on a cell that interacts with another molecule or macromolecule to confer a response or transduce a signal and includes nuclear receptors, mitochondrial receptors, cytoplasmic receptors, and cell surface receptors. Receptors include receptors for neurotrophins (including, without
5 limitation, nerve growth factor, neurotrophins 3, 4, and/or 5 (NT-3, NT-4 and/or NT-5) and brain derived growth factors); neurotransmitters; hormones; steroids; local mediators such as nitric oxide, carbon monoxide, histamine, and growth factors like insulin, insulin-like growth factor-I, fibroblast growth factors, ciliary neurotrophic factor, glia-derived neurotrophic factor, glia-derived nexin, cholinergic enhancing factor, transforming growth factors, activity
10 dependent neurotrophic factor, neurokines, gangliosides, phosphatidylserine, PDGF (platelet derived growth factor) and EGF (epidermal growth factor); benzodiazepines; arachidonic acid; purines (including, without limitation, adenosine and ATP); nucleotides and cyclic nucleotides; calcium and other divalent cations; odorants; antisense oligonucleotides; opiates; cannabinoids; glutamate; melatonin; angiotensin II; secretin; vasoactive intestinal peptide;
15 cholecystikinin; ACTH; vasopressin; thrombin; ion channels; and the like. Receptors also include but are not limited to G-protein-coupled receptors, ion-channel-linked receptors and enzyme-linked receptors.

As used herein, "protecting a receptor" refers to protecting the physical integrity of a receptor and/or the function of a receptor, such as enhancing the function of a receptor; or
20 maintaining the ability of the receptor to respond to agonists, to respond to antagonists, to transmit a message to the interior of a cell, or to send a signal within a cell, cell nucleus, or mitochondria.

As used herein, “central nervous system” (CNS) refers to the brain and spinal cord and associated tissues.

As used herein, “disorders and diseases of the CNS” refers to brain diseases such as Alzheimer’s disease, Parkinson’s disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, and/or schizophrenia; cell damage; nerve damage from cerebrovascular disorders such as stroke in the brain or spinal cord, from CNS infections including meningitis and HIV, from tumors of the brain and spinal cord, prion diseases, and CNS disorders resulting from ordinary aging (e.g., anosmia), brain injury, or spinal cord injury.

As used herein, a disease or disorder that relates to or is caused at least in part by dysfunction, alteration, or loss of one or more G-protein coupled receptors refers to Alzheimer’s disease; Parkinson’s disease; drug addiction, such as opiate addiction or cannabinoid abuse; pain; Sjogren’s or dry eye syndrome; heart diseases including congestive heart failure, myocardial infarction, cardiac arrhythmia; diseases of smooth muscle organs or glands such as irritable bowel syndrome, colitis, hypertension, erectile dysfunction, diabetes, obesity, blood coagulation disorders; and the like.

An “effective amount” of agent is an amount sufficient to prevent, treat, reduce and/or ameliorate the symptoms and/or underlying causes of any of the above disorders or diseases. In some instances, an “effective amount” is sufficient to eliminate the symptoms of those diseases and, perhaps, overcome the disease itself. Preferably, an effective amount of an agent yields a tissue concentration in the range of about 10^{-7} molar to about 10^{-5} molar, but the concentrations may be greater provided that toxicity is avoided.

In the context of the present invention, the terms “treat” and “therapy” and the like refer to alleviate, slow the progression, prophylaxis, attenuation or cure of existing disease. Prevent, as used herein, refers to putting off, delaying, slowing, inhibiting, or otherwise stopping, reducing or ameliorating the onset of such diseases or disorders. It is preferred that
5 a large enough quantity of the agent be applied in non-toxic levels in order to provide an effective level of activity against the disease. The method of the present invention may be used with any animal, such as a mammal or a bird (avian), more preferably a mammal. Poultry are a preferred bird. Exemplary mammals include, but are not limited to rats, cats, dogs, horses, cows, sheep, pigs, and more preferably humans.

Protecting a Tissue Component

The invention provides a method for protecting any biomolecule or tissue component including a protein, a lipid, a nucleic acid, a carbohydrate, a hormone and the like. The invention is best illustrated, but is not limited to, the example of the protection of a receptor,
15 preferably a muscarinic receptor, preferably a mAChR. In a first embodiment, the method includes administering a pyrophosphate analog. In a second embodiment, the method includes administering a protective agent. Protecting a receptor includes protecting the physical integrity of a receptor and/or the function of a receptor, such as maintaining the ability of the receptor to respond to agonists, to respond to antagonists, to transmit a message
20 to the interior of a cell, or to send a signal within a cell, cell nucleus, or mitochondria.

An embodiment of the invention provides a method for protecting a receptor from free radical damage. Free radicals and other reactive oxygen species (e.g., H_2O_2 , HOCl , and radicals such as O_2^- , sulfur cation, nitric oxide radical, ferryl, peroxy, peroxyxynitrite, thiyl,

thiylperoxyl, and alkoxy) are highly reactive, and many free radical reactions are highly damaging to cellular components. Free radical reactions can crosslink proteins, mutagenize DNA, and peroxidize lipids. Such reactions can have deleterious effects on cellular receptors. Preferably, the method of the invention includes protection of a receptor, such as a mAChR, or of DNA, RNA, lipids, and proteins necessary for receptor function from deleterious effects.

In another embodiment, the invention provides a method for reducing or eliminating deleterious effects of an endogenous inhibitor found in elevated levels in the brains of Alzheimer's disease patients. This endogenous, low molecular weight inhibitor, as it is known, inhibits agonist and antagonist binding to mAChRs. This inhibitor has a molecular weight of less than 3500 Da and is believed to generate free radicals, in the presence of glutathione or other sulthydryl compounds, that irreversibly inhibit or inactivate the mAChR. The inhibitor also contains free heme, which can generate free radicals, including superoxide radicals, peroxyl radicals, and thiyl radicals, and can cause neurotoxicity. Heme has been shown to damage protein and lipid components of membranes by Vincent (Oxidative Effects of Heme and Porphyrins on Proteins and Lipids, Seminars in Hematology 26(2): 105-113, 1989). Membrane lipid defects have been demonstrated in Alzheimer's disease by Ginsberg et al. (Evidence for a Membrane Lipid Defect in Alzheimer's Disease, Mol. and Chem. Neuropathol. 19: 37-46, 1993). In addition, heme has been proposed to contribute to atherosclerosis by Jacob (Newly recognized causes of atherosclerosis: The role of microorganisms and of vascular iron overload, J. Lab. Clin. Med. 123: 808-816, 1994).

In one embodiment, the method of the invention includes increasing the efficacy of an agent that directly or indirectly affects a mAChR. By way of example, administration of a pyrophosphate analog can increase the efficacy of a muscarinic agonist in the presence of the

inhibitor. In another embodiment, the method of the invention includes reducing or eliminating deleterious effects the low molecular weight inhibitor or heme by decreasing or preventing the generation of free radicals or trapping radicals once formed.

5 **Receptors**

The invention provides a method for protecting a receptor and/or increasing the efficacy of agents that directly or indirectly affect a receptor. Such receptors include G-protein-coupled receptors, ion-channel-linked receptors and enzyme-linked receptors.

10 Examples include receptors for neurotrophins; neurotransmitters; hormones; steroids; local mediators such as nitric oxide, histamine, and growth factors like PDGF (platelet derived growth factor) and EGF (epithelial growth factor); nucleotides and cyclic nucleotides; calcium and other divalent cations; odorants; antisense oligonucleotides; and the like. Preferably the receptor is a muscarinic receptor. Examples of G-protein coupled receptors include receptors that respond to odorants, opiates, cannabinoids, glutamate, melatonin, angiotensin II, secretin,
15 vasoactive intestinal peptide (VIP), cholecystokinin (CCK), adrenaline (adrenergic receptors), acetylcholine (muscarinic receptors), ACTH, vasopressin, thrombin, and the like.

In one embodiment the invention provides a method for protecting a receptor and/or increasing the efficacy of agents that directly or indirectly affect a receptor in the CNS. In another embodiment, the invention provides a method for protecting a receptor and/or
20 increasing the efficacy of agents that directly or indirectly affect a receptor not in the CNS. Agents whose efficacy are increased by the method of the invention include receptor agonists, allosteric modifiers of receptors, and receptor antagonists.

An embodiment of the invention provides a method for treating or preventing a disease or disorder that relates to or is caused at least in part by dysfunction, alteration, or loss of one or more G-protein coupled receptors. These diseases and disorders include Alzheimer's disease; Parkinson's disease; stroke; multiple sclerosis; ALS; drug addiction, such as opiate addiction or cannabinoid abuse; pain; Sjogren's or dry eye syndrome; heart diseases including congestive heart failure, myocardial infarction, cardiac arrhythmia; cancer; diseases of smooth muscle organs or glands such as irritable bowel syndrome, colitis, hypertension, erectile dysfunction, diabetes, obesity, blood coagulation disorders; and the like.

10 **Muscarinic Receptors**

The invention provides a method for protecting a mAChR and/or increasing the efficacy of agents that directly or indirectly affect a mAChR. There are at least five pharmacological classes of mAChRs, including the M1, M2, and M3 muscarinic receptors, and several genetic subclasses including m1, m2, m3, m4, and m5. These muscarinic receptors are G-protein coupled receptors. Each receptor subtype has its own unique pattern of expression throughout various tissues. As such, dysfunction of each receptor subclass, or combinations thereof, may have deleterious effects leading to a variety of diseases or disorders. The method of the invention can provide protection to a muscarinic receptor in any or several mAChR subclasses, and therefore, can be of benefit to those at risk or suffering from diseases associated with dysfunction of one or more muscarinic receptor subtype. Preferably, the method of the invention provides protection to M1 and M2 muscarinic receptors.

Muscarinic receptors mediate numerous of the inhibitory and excitatory effects of the neurotransmitter acetylcholine in the heart, smooth muscle, blood vessels, glands and in neurons (both presynaptic and postsynaptic) in the autonomic and the central nervous system. Dysfunction of mAChRs thus can contribute to a variety of diseases and/or disorders.

- 5 Through protection of a mAChR and/or through increasing the efficacy of agents that directly or indirectly affect a mAChR, the method of the invention can provide benefit to subjects suffering from or at risk of a disease or disorder associated with mAChR dysfunction.

- 10 An embodiment of the invention provides a method for protecting a mAChR and/or increasing the efficacy of agents that directly or indirectly affect a mAChR in the nervous system of a subject, and therefore, can be of benefit to subjects suffering from or at risk of central nervous system or peripheral nervous system disorders. For example, mAChRs and other receptors are involved in the regulation of the function of cells throughout the CNS. Accordingly, the method of the invention can provide benefit to subjects suffering from or at risk of CNS disorders such as Alzheimer's disease, Parkinson's disease, Lewy body dementia, multiple sclerosis, cerebellar ataxia, progressive supranuclear palsy, amyotrophic lateral sclerosis, affective disorders, anxiety disorders, and/or schizophrenia; nerve damage from cerebrovascular disorders such as stroke, from CNS infections including meningitis and HIV, from tumors of the brain and spinal cord, prion diseases, and CNS disorders resulting from ordinary aging, brain injury, or spinal cord injury. mAChRs are also involved in the regulation of the function of cells in the peripheral nervous system, including the autonomic nervous system. Accordingly, the method of the invention can provide benefit to subjects suffering from or at risk of peripheral nervous system disorders, such as peripheral neuropathy, including that associated with diabetes. For example, in diabetic patients nerves
- 15
- 20

can deteriorate as blood vessels that contain muscarinic or other receptors are lost.

Preferably, the method of the invention can benefit a subject suffering from or at risk of Alzheimer's disease.

Another embodiment of the invention provides a method for protecting a mAChR
5 and/or increasing the efficacy of agents that directly or indirectly affect a mAChR not within
the nervous system of a subject, and therefore, can be of benefit to subjects suffering from or
at risk of disease or disorder outside the nervous system. For example, mAChRs are involved
in the regulation (e.g., stimulation or inhibition) of smooth muscle contraction, the regulation
of heart rate and cardiac contractility, the regulation of secretion of enzymes or hormones,
10 including the release of amylase from the parotid gland and the release of digestive enzymes
and insulin from the pancreas, the regulation of bone growth, and the regulation of iron
metabolism. Accordingly, the method of the invention can provide benefit to subjects
suffering from or at risk of smooth muscle related disorders such as irritable bowel syndrome,
diverticular disease, urinary incontinence, esophageal achalasia, diseases or disorders of the
15 blood vessels (e.g. hypertension), or chronic obstructive airways disease; heart muscle related
disorders such as pathologic bradycardia or tachycardia, arrhythmia, flutter or fibrillation;
blood related disorders such as hereditary hemochromatosis; bone disorders such as Paget's
disease; cancer; and gland related disorders such as xerostomia, diabetes mellitus, or
Sjogren's syndrome or dry eye syndrome which involves decreased secretion of tears by, for
20 example, the lacrimal glands. For example, tear secretion is known to require muscarinic
cholinergic stimulation and intact muscarinic receptors. Accordingly, protection of a mAChR
and/or increasing the efficacy of an agent that directly or indirectly affects a mAChR can be
of benefit to a subject suffering from Sjogren's syndrome or dry eye syndrome.

Protecting a muscarinic receptor can result in enhancing the effectiveness of agents that directly or indirectly affect a mAChR. Useful agents that affect a mAChR include, but are not limited to, anticholinesterase agents, muscarinic agonists, muscarinic antagonists, and other agents useful for treatment of diseases associated with dysfunction of muscarinic
5 receptors, including neurodegenerative and other CNS disorders.

The method of the invention also provides enhanced efficacy of agents that do not act directly, or indirectly, with a mAChR. Such enhanced efficacy can be achieved, for example, through protection of a receptor, preferably a mAChR. Protecting a muscarinic receptor can result in enhancing the efficacy of agents that do not exert their action directly or indirectly on
10 the muscarinic receptor. Such enhanced efficacy can be achieved through desirable effects on cells to which protection of muscarinic receptors provides benefit. Typically, cells that derive benefit from protection of a muscarinic receptor are cells that contain muscarinic receptors. Examples of cells that contain a mAChR include particular neurons, smooth muscle cells, and gland cells.

15 Cells that lack a muscarinic receptor but which interact with enzymes, hormones, and/or other compounds released from cells with a muscarinic receptor can derive benefit from protecting a muscarinic receptor. Examples of cells lacking a mAChR that can derive benefit from protecting a mAChR include cells that are presynaptic or postsynaptic relative to cells that contain a mAChR and cells that can interact with enzymes, hormones, and/or other
20 compounds released from cells that contain a mAChR.

Examples of this phenomenon include: Stimulation of m_2 receptors on presynaptic membranes increases the release of acetylcholine which can then stimulate nicotinic receptors on other post synaptic cells. Stimulation of mAChR releases arachidonic acid, which can then

affect a variety of other nearby brain cells. Arachidonic acid also increases secretion of amyloid precursor protein. Emmerling, M.R. et al. (1996) *Ann. N.Y. Acad. Sci.* 777:310-315. Activation of m1 and m3 mAChR attenuates release of amyloid B protein. Hung, A.Y. et al. (1993) *J. Biol. Chem.* 268:22959-22962. Stimulation of mAChR is required for
5 memory and learning which also involve the proper function of noncholinergic cells. Stimulation of mAChR can increase the nitric oxide - cyclic GMP signaling system in neurons (Bauer, M.B. (1994) *Neuroscience* 62:351-359) and nitric oxide can travel from one cell to another to produce its effects. Stimulation of mAChRs markedly increase hippocampal BDNF and NGF in RNA levels. Once produced, these neurotrophins can produce important
10 effects on other nerve cells in the brain. M. da Penha Berzaghi (1993) *J. Neuroscience* 13(9) 3818-3826.

By way of further example, some cells containing a mAChR in the pancreas can release insulin. Released insulin can then interact with cells in close proximity to, or at relatively great distances from, the cell from which it was released. Protection of a mAChR
15 on a cell in the pancreas that releases insulin can have beneficial effects on a cell that interacts with insulin. Thus, cells lacking a mAChR can benefit from protection of a mAChR. Similarly, the efficacy of agents acting on cells lacking a mAChR can be enhanced by the method of the invention.


By way of yet further example, stimulation of mAChR in certain brain cells increases
20 potassium ion evoked release of the neurotransmitter dopamine which then goes on to affect other brain cells having dopamine receptors. Joseph, J.A. et al. (1995) *Brain Res.* 673:195-193. Thus, CNS cells lacking a mAChR can benefit from protection of a mAChR. Similarly,

the efficacy of agents acting on CNS cells lacking a mAChR can be enhanced by the method of the invention.

Pyrophosphate Analogs

- 5 In one embodiment, the method of the invention provides protection to a receptor and/or increases the efficacy of agents by administering to a subject a pyrophosphate analog.

Useful pyrophosphate analogs include compounds of Formula I:




- where each X is independently O, CH₂, NH, or S; R¹ is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²), and m is 1-3; R² is H, a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl; and n is 1-900. Compounds of Formula I in which R¹ is a small alkyl group, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, arachidonyl, -PO(OH)(OR²), or -(PO(OH)O)_m-PO(OH)(OR²); or R² is H, guanyl, adenylyl, glycerol, acyl glycerol, diacyl glycerol, serine, threonine, tyrosine, or arachidonyl can be referred to as substituted pyrophosphate analogs. Compounds of Formula I can also include substituted pyrophosphate analogs such as dinucleoside-5-5'-pyrophosphates, cyclophosphates of purine and pyrimidine acyclonucleosides. The compound of Formula I can be any pharmaceutically acceptable salt or basic addition salt. Preferably, X is O, CH₂, NH, or S; R¹ is H; and n is 2-6. More preferably the pyrophosphate analog is pyrophosphate or imidodiphosphate.

Additional preferred compounds of Formula I include pyrophosphate, glycerol pyrophosphate, arachidonylpyrophosphate, imidodiphosphate, serine phosphate, serine imidodiphosphate, threonine phosphate, threonine imidodiphosphate, guanylimidodiphosphate and adenylylimidodiphosphate. More preferably compounds of Formula I include

5 pyrophosphate, imidodiphosphate, guanylimidodiphosphate and adenylylimidodiphosphate.

Additional pyrophosphate analogs include compounds of formula II:



10 where $n=2-4$; X is O; RCR^1 ; CR; C ($n=4$), CH ($n=3$), or CH_2 ($n=3$); NH; N; S; and R and/or R^1 is H, OH, a small alkyl group (such as CH_3), or $(\text{CH}_2)_m\text{NH}_2$ where $m=1-6$. Further included are bisphosphonic acids, which are also known as bisphosphonates, where X is preferably RCR^1 and R and R^1 groups are chosen independently from OH, $\text{H}_2\text{N}(\text{CH}_2)_2$, or CH_3 . For example, RCR^1 can be $\text{H}_2\text{N}(\text{CH}_2)_2\text{C}(\text{OH})$ or CH_3COH . More specifically, the

15 bisphosphonates include etidronic acid ((1-Hydroxyethylidene)bisphosphonic acid) and pamidronic acid ((3-Amino-1-hydroxypropylidene)bisphosphonic acid) where preferably $n=2$.

Yet more additional pyrophosphate analogs include substituted pyrophosphate analogs such as inositol diphosphate, inositol triphosphate, inositol tetraphosphate, inositol pentaphosphate, and inositol hexaphosphate.

20 Such pyrophosphate and imidodiphosphate compounds and the like can be prepared as basic addition salts, such as sodium, potassium, or magnesium salts. It is believed that the use of a basic addition salt, such as a magnesium salt, will reduce the charge and allow for freer movement of the compound throughout the body. Pyrophosphate compounds,

imidopyrophosphate compounds, and the like can be covalently bound to other phosphates creating polyphosphates or polyimidophosphates. One or more pyrophosphate analogs can be administered in combination. In another embodiment, the pyrophosphate analog can be administered with a protective agent. In another embodiment, the pyrophosphate analog can

5 be administered with a neurologic agent, and optionally with a protective agent.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228

Protective Agents

In another embodiment, the invention provides a method for protecting receptors and/or increasing the efficacy of agents by administering to a subject a protective agent.

- 5 Protective agents useful in an embodiment of the method of the invention include a bilirubin, biliverdin, carnosol, quercetin, myricetin, a bioflavonoid; a heme binding compound, such as hemopexin, lipopexin, a lipoprotein, or ApoE-2; and a heme oxygenase, such as heme oxygenase-1 or heme oxygenase-2, or biliverdin reductase, a catalase, a peroxidase, a DNA or RNA vector encoding a biliverdin reductase, a DNA or RNA vector encoding a heme
- 10 oxygenase (e.g. a DNA or RNA vector encoding a heme oxygenase-1 or a DNA or RNA vector encoding a heme oxygenase-2), a DNA or RNA vector encoding a catalase, a DNA or RNA vector encoding a peroxidase, or a combination thereof Biliverdin reductase is preferably administered with bilirubin, because biliverdin reductase can regenerate bilirubin from biliverdin after bilirubin has been oxidized while functioning as a protective agent.
- 15 Biliverdin reductase is also preferably administered in combination with a heme oxygenase. Heme oxygenases include recombinant heme oxygenase. Preferably, a heme oxygenase is a human heme oxygenase.

One or more protective agents can be administered in combination. In another embodiment, one or more protective agents can be administered in combination with one or

20 more pyrophosphate analogs. In another embodiment, one or more protective agents can be administered in with one or more neurologic agents, and optionally with one or more pyrophosphate analogs.

Agents that Directly or Indirectly Affect a mAChR

The invention also provides a method for enhancing the efficacy of one or more agents that directly or indirectly affect a mAChR. Agents that directly or indirectly affect a mAChR include agents that (1) bind to or interact with a mAChR to either elicit a response transduced through a mAChR or reduce or prevent an agent from binding to or interacting with a mAChR and/or eliciting a signal transduced through a mAChR, (2) alter the concentration of agents that bind to or interact with a mAChR to either elicit a response transduced through a mAChR or reduce or prevent an agent from binding to or interacting with a mAChR and/or eliciting a signal transduced through a mAChR, or (3) modify the ability of agents that bind to or interact with a mAChR to either elicit a response transduced through a mAChR or reduce or prevent an agent from binding to or interacting with a mAChR and/or eliciting a signal transduced through a mAChR. Such agents include anticholinesterase agents, muscarinic agonists, muscarinic antagonists, and allosteric modifiers of muscarinic receptors. Preferably, the invention provides a method for enhancing the efficacy of agents that either directly or indirectly elicit a response through a mAChR. Most preferably, the method of the invention enhances the efficacy of a muscarinic agonist or an anticholinesterase agent.

Muscarinic receptor agonists directly elicit a response through a mAChR by binding to and transducing a signal through a mAChR. Preferred muscarinic agonists include acetylcholine, Xanomeline, and the like.

A muscarinic antagonist is an agent that is capable of partially or completely inhibiting, or reversing, the effect of a muscarinic agonist on a mAChR. Examples of muscarinic antagonists include atropine, N-methyl-scopolamine, quinuclidinyl benzilate, pirenzepine, and the like.

Cholinesterase hydrolyzes the neurotransmitter acetylcholine and provides one of the mechanisms responsible for rapid depletion of acetylcholine from the synaptic cleft.

Anticholinesterase agents inhibit cholinesterase activity and as a result increase the concentration of acetylcholine in the synaptic cleft and prolong the duration of which

5 acetylcholine remains in the synaptic cleft. Anticholinesterase agents can thus indirectly affect a mAChR by increasing concentrations of and prolonging the effective duration of acetylcholine in the synaptic cleft. Anticholinesterase agents can also interact directly with cholinergic receptors, including mAChRs; with sodium and potassium ion channels; and effect the uptake, synthesis and release of neurotransmitters. Preferred anticholinesterase
10 agents include Aricept, Exelon, Metrifonate, and the like.

An allosteric modifier of a mAChR binds or interacts with a site other than the agonist binding site of a mAChR and modifies the ability of an agonist or an antagonist to elicit or inhibit, respectively, a response transduced through a muscarinic receptor, without itself eliciting a response. Suitable allosteric modifiers of a muscarinic agonist include gallamine
15 and dynorphin. Preferred allosteric modifiers of a mAChR include gallamine and dynorphin.

Neurologic Agents

The invention also provides a method for increasing the efficacy of a neurologic agent in a subject in need thereof In a first embodiment, this method includes administering a
20 pyrophosphate analog. In this embodiment, the increased efficacy of the neurologic agent preferably results from protection of a muscarinic receptor caused or induced by the pyrophosphate analog. In a second embodiment, this method includes administering a protective agent. In this second embodiment, the increased efficacy of the neurologic agent

preferably results from protection of a muscarinic receptor caused or induced by the protective agent. In each embodiment, the subject preferably is concurrently receiving, has recently received, or will soon receive a neurologic agent.

A neurologic agent promotes nerve cell growth and survival or augments the activity of functioning cells. Among those agents that are preferred are cholinergic agonists, allosteric modifiers of a mAChR, cholinesterase inhibitors, or neurotrophic and neuritogenic factors that are similar to naturally occurring nerve growth promoting substances. Among the preferred neurologic agents are gangliosides (such as GM-I ganglioside), phosphatidylserine (PS), nerve growth factor (NGF), neurotrophins 3, 4, and/or 5 (NT-3, NT-4 and/or NT-5) brain-derived neurotrophic factor (BDNF), fibroblast growth factors (FGFs, e.g., basic fibroblast growth factor), insulin, insulin-like growth factors (IGF-1 and/or IGF-2), ciliary neurotrophic factor (CNTF), transforming growth factors, epidermal growth factors, activity-dependent growth factor, platelet derived growth factor, neurokinin, glia-derived neurotrophic factor (GDNF), glia-derived nexin, and cholinergic enhancing factors such as phosphoethanolamine and thyroid hormone T₃, and DNA or RNA vectors or plasmids that encode one or more protein neurologic agents or nerve growth promoting factors. Plasmids and vectors for delivery of a coding sequence to a mammalian tissue are known to those of skill in the art.

Metal Diseases

The method of the invention can treat or prevent diseases or disorders caused or induced by metals, such as cancer and poisoning. Such metals can include As, Co, Cr, Ni, Hg, Pb, Fe, Cu, V, and Cd. That is, the method of the invention can treat or prevent poisoning by (for example) lead or mercury and also excessive iron toxicity. In one embodiment, the

method of the invention can treat or prevent CNS diseases or disorders caused or induced by metals. In another embodiment, the method of the invention can treat or prevent diseases or disorders not of the CNS but caused or induced by metals, such as heart disease, blood vessel disease, and gland disease. In another embodiment, the method of the invention reduces poisoning of a subject by at least one metal. In another embodiment, the method of the invention protects a subject from at least one carcinogenic metal. In another embodiment, the method of the invention reduces toxic actions of metal ions in a subject, particularly toxic actions due to Fe^{++} , Hg^{++} , Cd^{++} , Cu^{++} , As^{+++} , and Pb^{++} ions.

10 Administering Agents

Administering compounds according to the method of the invention can include formulating the compounds or compositions as pharmaceutical compositions and administering the pharmaceutical compositions to a mammalian host, including a human patient, in a variety of forms adapted to the chosen route of administration. The compounds are preferably administered in combination with a pharmaceutically acceptable carrier. The compounds can be administered at one of a variety of doses sufficient to provide an effective amount at the desired point of action of the agent. Doses for humans and other mammals can range from about 0.001 mg/kg to about 100 mg/kg, preferably from about 0.01 mg/kg to about 10 mg/kg, preferably from about 0.1 mg/kg to about 1-10 mg/kg.

20 A related use of the methods of the invention is to protect pharmacological agents in formulation. The pharmacological agents may be for therapeutic, diagnostic, or other purposes.

The compounds can be administered by known techniques, such as orally, intranasally, parentally (including subcutaneous injection, intravenous, intramuscular, intrasternal or infusion techniques), by inhalation spray, dermally, transdermally, intrathecal, intracerebroventricular, buccal, sublingual, topically, by absorption through a mucous
5 membrane or through the skin, or rectally, in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants or vehicles.

Pharmaceutical compositions of the invention can be in the form of suspensions or tablets suitable for oral administration, nasal sprays, eye drops, nose drops, creams, sterile injectable preparations, such as sterile injectable aqueous or oleagenous suspensions or suppositories.

10 Controlled or sustained release systems can also be employed. For example, compositions can include a polymer or other substance that enhances controlled or sustained release. Controlled or sustained release systems can include a polymer disk, such as evac disks, microspheres, and copolymers. Preferred controlled release polymers are poly(lactide:glycolide) and poly(ethylene-co-vinyl acetate).

15 For oral administration as a suspension, the compositions can be prepared according to techniques well known in the art of pharmaceutical formulation. The compositions can contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents. As immediate release tablets, the compositions can contain microcrystalline cellulose,
20 starch, magnesium stearate and lactose or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

In addition to the typical pharmacological methods for oral administration, the agents employed in the methods of the invention can be administered as a component of a nutritional

or food supplement. The nutritional or food supplement can also include any other ingredients typical of a nutritional or food supplement, such as flavorings, stabilizers, and the like.

For administration by inhalation or aerosol, the compositions can be prepared according to techniques well known in the art of pharmaceutical formulation. The compositions can be prepared as solutions in saline, using benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons or other solubilizing or dispersing agents known in the art.

For administration as injectable solutions or suspensions, the compositions can be formulated according to techniques well-known in the art, using suitable dispersing or wetting and suspending agents, such as sterile oils, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

For intranasal administration, the compositions can be formulated according to techniques well known in the art. The means of applying a pharmaceutical composition intranasally can be in a variety of forms such as a powder, spray or nose drops.

For transdermal administration, the compositions can be formulated according to techniques well known in the art. Delivery of the composition through the skin can be accomplished by delivery means well known in the art, including transdermal patch, an ointment, an iontophoretic patch or device, and the like.

For rectal administration as suppositories, the compositions can be prepared by mixing with a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ambient temperatures, but liquefy or dissolve in the rectal cavity to release the drug.

Preferred administration routes include orally, parenterally, as well as intravenous, intramuscular or subcutaneous routes.

Intraocular administration through the use of an ointment or eye drops is preferred for treatment of a glandular disease or disorder of the eye, such as a lachrymal gland disease disorder like Sjogren's Syndrome or dry eye syndrome.

Solutions or suspensions of the compounds can be prepared in water, isotonic saline (PBS) and optionally mixed with a nontoxic surfactant. Dispersions may also be prepared in glycerol, liquid polyethylene, glycols, DNA, vegetable oils, triacetin and mixtures thereof Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

The pharmaceutical dosage form suitable for injection or infusion use can include sterile, aqueous solutions or dispersions or sterile powders including an active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium including, for example, water, ethanol, a polyol such as glycerol, propylene glycol, or liquid polyethylene glycols and the like, vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size, in the case of dispersion, or by the use of nontoxic surfactants. The prevention of the action of microorganisms can be accomplished by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be desirable to include isotonic agents, for example, sugars, buffers, or

sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the inclusion in the composition of agents delaying absorption--for example, aluminum monostearate hydrogels and gelatin.

Sterile injectable solutions are prepared by incorporating the compounds in the required amount in the appropriate solvent with various other ingredients as enumerated above and, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

Administering Agents to the Brain

Administering agents, e.g. a protective agent, a pyrophosphate analog, an agent that directly or indirectly affects a mAChR, and/or a neurologic agent, according to the method of the invention includes administering agents to a mammalian host in a manner that allows the agents to exert their effect in the CNS. Many agents useful for the method of the invention can be absorbed into the blood stream and readily cross the blood brain barrier.

However, some agents useful for the method of the invention cannot pass, or have difficulty passing, the blood brain barrier. Such agents can be administered as "prodrugs" which can cross the blood brain barrier, and upon or after entry into the CNS, the prodrug is converted to the active agent. Agents that can cross the blood brain barrier without difficulty can also be administered as prodrugs.

To deliver the agent to the CNS, the agent alone or in combination with other substances as a pharmaceutical composition may be administered to the spinal cord and to the

cerebral vesicles according to intrathecal and intracerebrovascular administration methods known in the art. Such pharmaceutical compositions can also be administered to the nasal cavity, under the tongue, or onto the eye. The composition may be dispensed intranasally, sublingually, or conjunctivally as a powdered or liquid nasal spray, nose drops, a gel or ointment, through a tube or catheter, by syringe, by packtail, by pledget, or by submucosal infusion. The agent may be combined with a polymer or other substance that enhances controlled or sustained release of the agent. In particular, agents can be delivered to the brain by intranasal administration as described in X.-Q. Chen et al. (1998) J. Alzheimer's Disease 1:35-44 and W.H. Frey II et al. (1997) Drug Delivery 4:87-92; the disclosures of which are incorporated herein by reference.

The optimal concentration of the active agent will necessarily depend upon the specific agent used, the characteristics of the patient and the nature of the disease or condition for which the treatment is to be used.

The carrier of the composition may be any material which is otherwise pharmaceutically-acceptable and compatible with the active ingredients of the composition. Where the carrier is a liquid, it is preferred that the carrier is hypotonic or isotonic with nasal, oral, or conjunctival fluids and have a pH within the range of 4.5-7.5. Where the carrier is in powdered form, it is preferred that the carrier is also within an acceptable non-toxic pH range.

The pharmaceutical composition may be formulated as a powder, granules, solution, ointment, cream, aerosol, powder, drops, or a controlled or sustained release composition such as a polymer disk. The solution may be sterile, isotonic or hypotonic, and otherwise suitable for administration by injection or other means. In addition to the agent, the solution may contain appropriate adjuvants, buffers, preservatives and salts. The powder or granular forms

of the pharmaceutical composition may be combined with a solution and with diluting, dispersing or surface active agents. Solutions such as nose or eye drops may contain an antioxidant, a buffer, and the like. Further controlled release polymers may be used to regulate the delivery of the agent.

- 5 The present invention may be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

EXAMPLES

10 **Example 1 - Protection of Muscarinic Acetylcholine Receptor (mAChR) in Cell Free Systems**

Materials and Methods

Membrane mAChR Preparation

- 15 Membranes rich in mAChRs were prepared by a modification of the method used by Marks and Collins (Characterization of nicotine binding in mouse brain and comparison with the binding of α -bungarotoxin and quinuclidinyl benzilate, Mol. Pharmacol. 22:544-564, 1982). Gray matter from nondemented adult human frontal cortex was homogenized in 9 vol of 50 mM Tris-HCl, pH 7.4, using 5 passes of a glass/Teflon motor-driven homogenizer. The homogenate was centrifuged at 27 000 x g for 20 min at 4°C, and the subsequent pellet
- 20 resuspended in 9 vol of cold deionized water with 5 passes of the homogenizer. The resuspension was incubated at 37°C for 5 min, then was centrifuged as before. The resulting pellet was resuspended, incubated and centrifuged again as above. The final pellet was weighed, resuspended at 15% w/v in 50 mM Tris-HCl buffer, aliquoted in small portions,

flash-frozen in liquid nitrogen and stored at -70°C for subsequent assays to determine content and binding capacity. Before use in binding assays, the thawed membrane preparation was briefly rehomogenized with 10 passes in a glass/glass homogenizer. A typical mAChR membrane preparation bound 300 pmol [³H] quinuclidinyl benzilate ([³H]QNB)/g protein.

5

Inhibitor Preparation

Gray matter obtained from the frontal cortex of cases with AD was homogenized in 9 vol of 1% trifluoroacetic acid (TFA) for 40 s at 4°C in a Waring blender, then centrifuged at 1200 x g for 10 min at 4°C. The resulting supernatant fraction was centrifuged at 11 000 x g for 100 min at 4°C. The 11 000 x g supernatant fraction was centrifuged at 100 000 x g for 100 min at 4°C, then the 100 000 x g supernatant fraction was concentrated using a SpeedVac and resuspended in 0.1% TFA to half the original tissue volume. The 100 000 x g supernatant fraction was transferred to a Spectra/Por 3 dialysis membrane bag (3500 dalton cutoff), and dialyzed against 20 vol of 0.1% TFA at 4°C for 24 h with gentle stirring. The resulting <3500 Da fraction (dialysate) was concentrated by SpeedVac to half the original tissue volume. The <3500 Da fraction (endogenous inhibitor) was then frozen in liquid nitrogen and stored at -70°C for subsequent assays to determine protein count and inhibitor activity. Protein activity was measured using the bicinchononic acid (BCA) protein assay method, essentially as described by Smith et al. (Measurement of protein binding using bicinchononic acid, Ann. Biochem. 150: 76-85, 1985). A typical inhibitor preparation contained about 4 mg/ml protein and approximately twice the concentration of inhibitor found in the original tissue.

Inhibitor Activity Assay

Inhibitor activity was measured using a modification of the method of Fields et al. (Cardiac muscarinic receptors, J. Biol. Chem. 253:3251-3258, 1978) to assess the binding of [³H]QNB, a mAChR antagonist, or [³H]-oxotremorine M, a mAChR agonist. In general, binding conditions consisted of 50 mM Tris-HCl, pH 7.4 at 37°C, 10 mM reduced glutathione (GSH), 75 µM membrane and 2 x 10⁻¹⁰ M [³H]QNB or 3nM [³H]-oxotremorine M, with and without addition of inhibitor. To control for non-specific binding, 12.5 µM atropine sulfate (a mAChR antagonist) was added to several tubes. Subtracting nonspecific binding from total binding yielded specific binding.

Pyrophosphate, imidodiphosphate, adenylylimidodiphosphate, guanylimidodiphosphate, and tripolyphosphate were dissolved in distilled water. Bilirubin, biliverdin, and heme were dissolved in DMSO. Carnosol, myricetin, and quercetin were dissolved in ethanol. Catalase and peroxidase were dissolved in an aqueous, preferably buffered, solution.

Enough water was added to all other reaction components in each tube to make 4 ml total. The binding reaction was initiated by adding [³H]QNB or [³H]-oxotremorine-M, mixing the tubes briefly, and then incubating the tubes at 37°C for [³H]QNB or at room temperature for [³H]-oxotremorine-M. The reaction time for [³H]QNB was one hour in most experiments. In some experiments, the mAChR was preincubated with either the endogenous LMW inhibitor or heme plus peroxide in the presence or absence of the therapeutic agent being tested. The effect of the therapeutic agent on receptor function was then assessed in a binding assay, which for [³H]QNB was conducted at 37°C for 40 min, and for [³H]-oxotremorine-M was conducted at room temperature for 20 min. After 60 min, the binding reaction was terminated by adding 5 ml of cold 50 mM Tris buffer, pH 7.4, to each tube and

chilling the tubes in an ice bath. The tube contents and one 15 ml rinse of cold 50 mM Tris buffer, pH 7.4, were filtered through Whatman GF/B glass fiber filters using a Brandel harvester. The filters were placed in Optiflow scintillation flour and counted in a Beckman LS-6500 scintillation counter set for tritium detection.

5

Results

The data resulting from the methods presented above and the results presented in Figures 1-20 are discussed in more detail below.

10 Pyrophosphate:

Pyrophosphate protects the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide. Pyrophosphate protected the receptor from both loss of antagonist (^3H -QNB) binding (Figures 1 and 2) and agonist (^3H -Oxotremorine M) binding (Figure 3). Approximately 1 μM pyrophosphate provides 50% protection. Pyrophosphate
15 also protects the mAChR from damage by PbCl_2 . Approximately 57 μM pyrophosphate provides 50% protection (Figure 20).

Imidodiphosphates:

Imidodiphosphate (Figure 4), guanylimidodiphosphate (Figure 5), and
20 adenylylimidodiphosphate (Figure 6) all protect the mAChR from inactivation by the LMW inhibitor.

Polyphosphates:

Polyphosphates, such as tripolyphosphate (Figure 7), protect the mAChR from inactivation by the LMW inhibitor.

Bisphosphonates:

- 5 Bisphosphonates, such as pamidronate (Figure 19), protect the mAChR from inactivation by the LMW inhibitor.

Bilirubin and Biliverdin:

- 10 Bilirubin protects the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide. Approximately 0.7 μ M bilirubin provides 50% protection of the receptor from loss of antagonist (3 H-QNB) binding (Figures 8 and 9) and 1.9 μ M provides 50% protection from loss of agonist (3 H-Oxotremorine M) binding (Figure 10).
 biliverdin at 3 μ M provides 50% protection of the mAChR (Figure 11).

15 Carnosol, Quercetin, and Myricetin:

Carnosol (Figure 12), quercetin (Figure 13), and myricetin (Figure 14) all protected the mAChR from inactivation. Carnosol provided 100% protection at 1 μ M. while quercetin and myricetin provided 50% protection at 0.24 μ M and 0.4 μ M respectively.

20 Catalase and Peroxidase

Catalase protected the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide (Figures 15 and 16, respectively). As little as 0.34 units/mL of catalase provided 50% protection from inactivation by heme and peroxide.

A peroxidase, specifically glutathione peroxidase, protected the mAChR from inactivation by the LMW inhibitor or by the combination of heme and peroxide (Figures 17 and 18, respectively). Glutathione peroxidase at 0.5 units/mL provided 71% protection from inactivation by heme and peroxide.

5

Conclusion

The results indicate that pyrophosphate, imidodiphosphates, polyphosphates, bisphosphonates, bilirubin, biliverdin, carnosol, quercetin and myricetin protect a receptor and increase the ability of agents to bind a receptor. Particularly, the results demonstrate the ability of these agents to protect a muscarinic receptor from the effects of endogenous LMW inhibitor, heme and metals and increase the ability of muscarinic agonists and antagonists to bind a mAChR, suggesting that these agents can be used effectively to protect other receptors and increase the efficacy of other agents.

Because the mAChR is essential for memory and learning, the specific demonstration that these agents can protect the human brain mAChR from inactivation and increase agonist binding indicates that these agents have therapeutic potential for the treatment of cognitive and memory disorders including those associated with aging, such as Alzheimer's disease.

Example 2- Protection of the mAChR in Cell Culture

Various systems for determining the protection of a mAChR in cell culture are known in the art. Such cell culture systems can be used to determine if mAChR is protected from a damaging agent or condition according to the method of the invention. For example, by administering a mAChR antagonist or agonist alone or in combination with one or more

protective agent and/or one or more pyrophosphate analog, one of skill in the art can determine if a mAChR is protected by the one or more protective agent and/or one or more pyrophosphate analog.

5 **Example 3- Protection of the mAChR Receptor in Animals**

Various systems for determining the protection of a mAChR in animals are known in the art. Such animal systems can be used to determine if mAChR is protected according to the method of the invention. For example, by administering a mAChR antagonist or agonist alone or in combination with one or more protective agents and/or one or more pyrophosphate
10 analogs, one of skill in the art can determine if a mAChR is protected by the one or more protective agents and/or one or more pyrophosphate analogs.

Example 4- Increased Efficacy of Neurologic Agents in Model Systems

Various models systems for determining the efficacy of neurologic agents are known
15 in the art. Such model systems can be used to determine if the efficacy of a neurologic agent is increased by the method of the invention. For example, by administering one or more neurologic agents alone or in combination with one or more protective agents and/or one or more pyrophosphate analogs, one of skill in the art can determine if the efficacy of the one or more neurologic agents is increased when administered with one or more protective agents
20 and/or one or more pyrophosphate analogs within the parameters of the model system according to techniques known in the art.

It should be noted that, as used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates

otherwise. Thus, for example, reference to a composition containing “a compound” includes a mixture of two or more compounds. It should also be noted that the term “or” is generally employed in its sense including “and/or” unless the content clearly dictates otherwise.

All publications and patent applications in this specification are indicative of the level
5 of ordinary skill in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated by reference.

The invention has been described with reference to various specific and preferred
10 embodiments and techniques. However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention.